

$$F_{\nabla} = 2\pi \cdot r^3 \frac{\sqrt{\epsilon_B}}{c} \left(\frac{\epsilon - \epsilon_B}{\epsilon + 2\epsilon_B} \right) (\nabla \cdot I)$$

F_{∇} = Optical force on particle towards higher intensity

r = Radius of particle




ϵ_B = Dielectric constant of background medium

ϵ = Dielectric constant of particle

I = Light intensity (W/cm^2)

∇ = Spatial derivative

FIG. 1



3/19




FIG. 4A




FIG. 5






FIG. 6

**FIG. 8**




FIG. 9A




FIG. 9B




FIG. 10




FIG. 11A



FIG. 11B




FIG. 11C

3916
100% 0.00
100% 0.00




FIG. 12A




FIG. 12B

FIG. 12C




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 14A




FIG. 14B




FIG. 15




FIG. 17




FIG. 16




FIG. 18




FIG. 20





FIG. 21

12/19





FIG. 22

FIG. 23






FIG. 24




FIG. 25

Before:

SCATTER FORCE SEPARATION



After:

FIG. 26

3910
SEARCHED
INDEXED
MAILED
JULY 1984
FBI LABORATORY

FIG. 27A




FIG. 27B




FIG. 27C




FIG. 28





FIG. 29






FIG. 30


HEMOGLOBIN - O₂ ABSORPTION SPECTRUM

FIG. 31





FIG. 32



FIG. 33



Before

After

Difference

FIG. 34

DISTRIBUTION OF ESCAPE VELOCITIES
 READING TAKEN IN PBS/1% BSA BUFFER
 RAIN-X COATED SLIDE/CYTOP COATED COVERSUP




FIG. 35




FIG. 36